

# **Proximal hamstring repair – surgical technique**

Lasse Lempainen

Hospital Mehiläinen NEO, Turku, Finland

Correspondence:

Lasse Lempainen, MD, PhD

Adjunct professor, orthopaedic surgeon

Hospital Mehiläinen NEO

Joukahaisenkatu 6

FI-20520 Turku, Finland

Email: [lasse.lempainen@utu.fi](mailto:lasse.lempainen@utu.fi), [lasse.lempainen@mehilainen.fi](mailto:lasse.lempainen@mehilainen.fi)

## **Abstract**

Proximal hamstring injuries are common in athletes. Surgical indications are based on to regain the pre-injury activity level of the athlete.

The key to achieve athletes back to play is to understand that hamstrings consist of three different muscles which have different functions.

The rupture of proximal hamstring tendon can occur to one, two or all of these three tendons (semitendinosus, semimembranosus and biceps femoris) which may result to disabling and chronic pain with unable to fully participate to competitive sports.

Surgery with open technique provides good visibility to hamstring tendons, nervous structures and bony insertion landmarks.

This chapter gives the reader the indications for proximal hamstring tendon avulsion surgery together with the surgical technique and post-operative rehabilitation. We also discuss possible complications of the operative treatment.

Keywords: hamstring injury, tendon avulsion, open surgery, surgical technique, rehabilitation

## **Introduction**


Hamstring injuries can occur in various sports activities but also among regular people eg while falling down. These injuries can be highly disabling and they can lead to substantial time loss from sports (Harris et al, 2011). Some of the hamstring injuries require surgical treatment for optimal recovery (Lempainen 2015). Various open surgery techniques have been presented in literature (Lempainen 2006, Pombo & Bradley, 2009; Cohen et al, 2012; Sarimo et al, 2008; Birmingham et al, 2011; Carmichael et al, 2009; Moatshe et al, 2017). The goal of the proximal hamstring rupture repair is to restore the anatomy of the injured structures so that athlete's rapid recovery and safe return sports is possible with the low rate of recurrent hamstring injuries. Without adequate treatment, proximal hamstring rupture can result in permanent loss of hamstring function and strength and also chronic pain (Shambaugh et al, 2017; Harris et al, 2011).

## **Surgical indications**


Most hamstring injuries are strains and can be treated conservatively with good results. However, there are cases in which surgery should be considered already in the acute phase. Also, there are cases in which surgery should be considered later if non-operative treatment appears to be unsuccessful.

The physician uses clinical findings (posterior thigh hematoma, pain and decreased strength in hip extension / knee flexion) and MRI imaging to determine whether the athlete has complete (Figure 1, 3) or incomplete proximal hamstring rupture (Figure 2, 4).

Figure 1. Complete proximal 3-tendon avulsion: A) axial; B) coronal; C) sagittal planes



**Figure 2. Complete proximal 1-tendon avulsion (semimembranosus): A) sagittal; B) axial; C) coronal (1) with subsequent axial view (2).**



#### *Absolute indications for surgery*

In an athlete a proximal one tendon avulsion / rupture with a clear retraction should be treated surgically regardless of the hamstring tendon (BF, SM or ST) (Figure 2, 4). If two or all three of the hamstring muscles are avulsed surgery should be considered in all patients if there are not contraindications to surgery. Suture anchors are typically used to reattach the tendon to the ischial tuberosity.

Apophyseal avulsions of the ischial tuberosity occur occasionally in adolescent athletes (Sinikumpu et al, 2018). Surgical repair is traditionally recommended if the avulsed fragment is displaced by more than 10-15 mm.

#### *Relative indications for surgery*

Occasionally incomplete tears especially when recurrent form scar tissue and adhesions that cause persistent symptoms and are non-responsive to conservative treatment. This can occur in the proximal interface or in the proximal tendinous part or in the central tendon area. In proximal incomplete avulsions that remain symptomatic the MRI may show liquid between the bone and the tendon which is a sign of incomplete healing.

It has been suggested that paramuscular / central tendon injuries especially in the BF may have a higher risk of poor healing with conservative treatment (Lempainen 2018). Also, the risk of a recurrent injury may be high. In these injuries there is often an incomplete tear of the paramuscular tendon typically in the area of 5 to 20 cm from the proximal origin. Often the muscle tissue is torn off from the tendon also. When a tear like this remains symptomatic after adequate conservative treatment or there are recurrences surgery should be considered. Full continuity of the central tendon is restored with sutures and the attachment of the muscle to the tendon is reinforced. It is important to avoid overtightening of the repaired tendon. Scar tissue may be removed. Suture anchors may be used if the tear is located close to the bony origin.


In chronic proximal hamstring ruptures and in some re-rupture cases, anatomic apposition of the retracted muscles cannot always be achieved. In those cases fascia lata autograft augmentation or e.g. Achilles allograft have been used to connect the retracted hamstrings to ischial tuberosity (Lempainen et al, 2007, Folsom & Larson, 2008). If patient has radiating nerve pain, sciatic nerve is typically adhered to the scar tissue and it should be freed for good outcome. It seems that late reconstruction of complete proximal hamstring avulsion with fascia lata autograft augmentation or allograft can result in enhancement of muscle strength, better function of the hamstrings and improved the leg control. Also symptoms derived from retracted hamstrings causing stretching to the sciatic nerve could be alleviated.

## Surgical technique

### Patient positioning and preparation

In surgery the patient is placed in the prone position and usually spinal anesthesia is used. The ipsilateral knee is slightly flexed (20°) to relax the hamstring muscles. The whole area of the hamstrings should be prepared especially in more chronic cases.

**Figure 3. Surgical approach and anatomy: A) Retracted complete proximal 3-tendon avulsion identified using vertical skin incision; B) Suture anchors placed anatomically to ischial tuberosity; C) Prepared and suture-loaded tendons are secured to the footprint.**




A vertical skin incision should be used especially when there is retraction of the ruptured tendons as they may need to be mobilized to achieve tension-free contact to the ischial tuberosity again. The incision starts at the ischial tuberosity extending distally, approximately 10-15 cm. A fasciotomy is done to the common hamstring fascia and it is continued distally, approximately 15

cm from the origin of the hamstring muscles. The lower edge of the gluteus maximus muscle is freed and careful hemostasis should be performed. The posterior cutaneous femoral nerve should be identified and spared as well. This is not always easy, especially in the chronic cases. The ischial tuberosity is exposed by retracting superiorly the inferior border of the gluteus maximus muscle. The sciatic nerve can be found lateral to the ischial tuberosity and it should be freed from adhesions in chronic cases, especially if there are sciatica type of symptoms.

In acute ruptures, hematoma or seroma is often present. In more chronic ruptures, adhesions and scarring is covering the ruptured area. In chronic cases ruptured structures should be carefully freed and mobilized like sciatic nerve.

**Figure 4. Retracted single tendon (biceps femoris) avulsion identified using vertical skin incision; B) Suture anchors placed anatomically to ischial tuberosity; C) Prepared and suture-loaded tendon is secured to the footprint.**



### *Surgical repair*

In most cases, re-attachment of the torn tendons can be done using suture anchors. When surgery is performed in the early phase, anatomical reattachment can be done. In chronic cases, the torn



tendons can be reattached slightly distal and medial to the original site of the ischial tuberosity to avoid over tightening the tendons.

Anchor placement to the ischial tuberosity should be anatomical and typically 2-3 suture anchors are used to reinsert ruptured tendons back to the bone. Suture passing through tendon should be done carefully and multiple times and then the ruptured tendon is reinserted to the footprint area by pulling the gliding strand of the suture anchor. Care must be taken not to rotate or misplace tendon heads. After the repaired proximal hamstring tendons are in good contact for prepared bony cortex surface, knots can be done.

If proximal part of the tendon / central tendon is ruptured the goal of the repair is to restore full continuity of the tendon with sutures and also the attachment of the muscle to the tendon is reinforced. It is important to avoid over tightening of the repaired tendon. Scar tissue may be removed. Suture anchors may be used if the central tendon tear is located close to the bony origin.

Wound closing is done by layers; subcutaneous tissue and skin. The skin sutures should be removed after 10 – 12 days.

## **Post-operative Rehabilitation Protocol**

- complete proximal hamstring rupture (acute repair, no augmentation)
- isolated proximal biceps femoris rupture / conjoint tendon (BF + ST) rupture
- proximal semimembranosus rupture

Postoperatively, no immobilization with casts or orthoses are needed. The patients are allowed to begin full weight bearing immediately after surgery. However, often crutches are needed 1-3 weeks after operation. Sitting is avoided for the first 2-3 postoperative weeks.

General principles of rehabilitation and routine protocol:

- Often daily surgery; patient can leave the hospital same day.
- First knee slightly flexed while laying on bed.
- Wound check after 2-3 days, sutures removed after 10-12 days.
- Walking with help of two crutches during first 1-3 weeks. However, full weight bearing is allowed immediately after operation while standing and slow walking.
- Calf and gluteus muscle activations can be started right after operation as well as isometric hamstring contractions.
- Sitting should be avoided as much as possible during first three weeks.
- Active stretching of the hamstrings should be avoided first four / five weeks.
- Functional strengthening / physiotherapy starts normally at 4 weeks. Gradually increasing load of the hamstrings. It is also important to concentrate on to the gluteus, calf muscles and pelvis core training.
- Light aqua training can be started after 3-4 weeks, cross-trainer / stationary biking after 5-6 weeks, Alter-G running after 8-10 weeks, normal running after 2-3 months and return to field after 2.5-4 months from the operation.
- Return to high level of sports after 3-5 months from the operation when pain-free and safe sports related movements are successfully performed.

## **Complications**

Severe complications are possible relating to the hamstring surgery. One should be aware and experienced about general principles of muscle tendon surgery. Some of these complications are related to the injury itself and some to the surgical technique.

There are some cases presented in the literature in which a complete proximal hamstring rupture has caused a dysfunction of the sciatic nerve with resulting complete foot drop and numbness of the lateral calf and dorsal foot (Street and Burks 2000, Hernesman et al. 2003). Similar cases have been described occurring after partial proximal hamstring tears (Carmody and Prietto 1995, Takami et al. 2000). The damage to the intramuscular nerve branches of the hamstring muscles is also a possible complication and could be seen e.g. in chronic proximal avulsion injuries. Nerve branch lesions can be also result from the surgery. If denervation of the hamstring muscles is suspected before operation ENMG study is recommended to do.

Lesion of posterior femoral cutaneous nerve is also possible and that is especially related to the cases needing reoperation. Most of these patients have some numbness around the scar area which do not cause any harm in their activities of normal daily life.

Post-operative infections are very rare in good planned and performed operations. We use antibiotic prophylaxis routinely before operation. After operation careful wound control(s) is of course important as well. Sutures should not be kept for too long time; 10 days is normally enough.

Good knowledge of the anatomy and 'experienced hands' are important and essential to avoid surgery related failures. Hamstring operations are often demanding procedures and should be

done by surgeons specialized for these injuries. One should be aware how to handle sciatic nerve, how to place and insert anchors to the ischial tuberosity and how to prepare tendons and take the suture attachments from the ruptured tendons.

## **Conclusion**

In competitive athlete, open surgical treatment for proximal hamstring tears gives good outcome (Lempainen et al, 2009, Lempainen 2009 Thesis). Indications for operative and non-operative treatment depends of the activity level of the patient, the retraction of the avulsion and the number of avulsed tendon heads (Lempainen et al 2015, Barnett et al, 2015; Cohen et al, 2012; Shambaugh et al 2017; Piposar et al, 2017). The vertical incision gives good visualization to the injured tendons and neural structures. By using modern suture anchors safe and tissue friendly reattachments are possible, ruptured tendons can be pulled to the anatomic insertion securely which enables the strength of the hamstrings be restored. After uneventful operation the athlete can return-to-play within a one competitive season if the rehabilitation is done by well-trained lower extremity specialized physiotherapist and staged rehabilitation program (Lempainen 2009, Lempainen et al 2018).

## References

- Barnett AJ, Negus JJ, Barton T, Wood DG. Reattachment of the proximal hamstring origin: outcome in patients with partial and complete tears. *Knee Surg Sport Traumatol Arthrosc.* 2015; 23:2130–5.
- Birmingham P, Muller M, Wickiewicz T, Cavanaugh J, Rodeo S, Warren R. Functional outcome after repair of proximal hamstring avulsions. *J Bone Joint Surg Am.* 2011;93:1819-26.
- Carmichael J, Packham I, Trikha SP, Wood DG. Avulsion of the proximal hamstring origin. Surgical technique. *J Bone Joint Surg Am.* 2009;91: Suppl 2:249-56.
- Carmody C, Prietto C. Entrapment of the sciatic nerve as a late sequela of injury to the hamstring muscles. A case report. *J Bone Joint Surg Am.* 1995;77:1100-2.
- Cohen SB, Rangavajjula A, Vyas D, Bradley JP. Functional results and outcomes after repair of proximal hamstring avulsions. *Am J Sports Med.* 2012;40:2092-8.
- Harris JD, Griesser MJ, Best TM, Ellis TJ. Treatment of proximal hamstring ruptures - a systematic review. *Int J Sports Med.* 2011;32:490-5.
- Hernesman SC, Hoch AZ, Vetter CS, Young CC. Foot drop in a marathon runner from chronic complete hamstring tear. *Clin J Sport Med.* 2003;13:365-8.
- Lempainen L, Sarimo J, Heikkilä J, Mattila K, Orava S. Surgical treatment of partial tears of the proximal origin of the hamstring muscles. *Br J Sports Med.* 2006;40:688-91.
- Lempainen L, Sarimo J, Orava S. Recurrent and chronic complete ruptures of the proximal origin of the hamstring muscles repaired with fascia lata autograft augmentation. *Arthroscopy.* 2007;23:441.e1-5.
- Folsom GJ, Larson CM. Surgical treatment of acute versus chronic complete proximal hamstring ruptures: results of a new allograft technique for chronic reconstructions. *Am J Sports Med.* 2008;36:104-9.
- Lempainen L, Sarimo J, Mattila K, Vaitinen S, Orava S. Proximal hamstring tendinopathy: Results of surgical management and histopathologic findings. *Am J Sports Med.* 2009;37:727–4.

Lempainen, L. Thesis. Surgical treatment of hamstring injuries and disorders – the clinical spectrum from chronic tendinopathy to complete rupture. University of Turku, Turku, Finland, 2009.  
<http://www.doria.fi/bitstream/handle/10024/43989/AnnalesD840Lempainen.pdf>

Lempainen L, Banke IJ, Johansson, Brucker PU, Sarimo J, Orava S, Imhoff AB. Clinical principles in the management of hamstring injuries. *Knee Surg Sports Traumatol Arthrosc.* 2015;23:2449-56.

Lempainen L, Kosola J, Pruna R, Puigdemivoll J, Sarimo J, Niemi P, Orava S. Central Tendon Injuries of Hamstring Muscles: Case Series of Operative Treatment. *Orthop J Sports Med.* 2018;6:2325967118755992.

Moatshe G, Chahla J, Vap AR, Ferrari M, Sanchez G, Mitchell JJ, LaPrade RF. Repair of Proximal Hamstring Tears: A Surgical Technique. *Arthrosc Tech.* 2017;6:e311–7.

Piposar JR, Vinod AV, Olsen JR, Lacerte E, Miller SL. High-Grade Partial and Retracted (<2 cm) Proximal Hamstring Ruptures: Nonsurgical Treatment Revisited. *Orthop J Sports Med.* 2017;5:2325967117692507

Pombo M, Bradley JP. Proximal Hamstring Avulsion Injuries - A Technique Note on Surgical Repairs. *Sports Health.* 2009;1:261–4.

Sarimo J, Lempainen L, Mattila K, Orava S. Complete proximal hamstring avulsions: a series of 41 patients with operative treatment. *Am J Sports Med.* 2008;36:1110-5.

Shambaugh BC, Olsen JR, Lacerte E, Kellum E, Miller SL. A Comparison of Nonoperative and Operative Treatment of Complete Proximal Hamstring Ruptures. *Orthop J Sports Med.* 2017;5:2325967117738551.

Sinikumpu JJ, Hetsroni I, Schilders E, Lempainen L, Serlo W, Orava S. Operative treatment of pelvic apophyseal avulsions in adolescent and young adult athletes: a follow-up study. *Eur J Orthop Surg Traumatol.* 2018;28:423-9.

Street CC, Burks RT. Chronic complete hamstring avulsion causing foot drop. A case report. *Am J Sports Med.* 2000;28:574-6.

Takami H, Takahashi S, Ando M. Late sciatic nerve palsy following avulsion of the biceps femoris muscle from the ischial tuberosity. *Arch Orthop Trauma Surg.* 2000;120:352-4.